

AD-A230 109

100 FILE 0011

2

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 34

Spectrum of the Third-Order Nonlinear Susceptibility of Trans-Polyacetylene

by

X. Sun, Z. Shuai, R. Fu, K. Nasu, X. S. Li, D. L. Lin and Thomas F. George

Prepared for publication

in

Journal of Physics: Condensed Matter (Letters)

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

December 1990

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
DEC 17 1990
S D
Ck

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/90/TR-34			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260				7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Spectrum of the Third-Order Nonlinear Susceptibility					
12. PERSONAL AUTHOR(S) X. Sun, Z. Shuai, R. Fu, K. Nasu, X. S. Li, D. L. Lin and Thomas F. George*					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) December 1990	
				15. PAGE COUNT 7	
16. SUPPLEMENTARY NOTATION Prepared for publication in <i>Journal of Physics: Condensed Matter</i> (Letters)					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	NONLINEAR SUSCEPTIBILITY TWO-PEAK SPECTRUM		
			THIRD-ORDER THEORY		
			TRANS-POLYACETYLENE FINITE CHAIN		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) By considering the finite lifetime of excitation and the band structure of a finite polymer chain, the two-peak structure in the spectrum of the third-order nonlinear susceptibility of trans-polyacetylene is calculated, with results in agreement with experiments.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Spectrum of the Third-Order Nonlinear Susceptibility

of Trans-Polyacetylene

X Sun^{††}, Z Shuai[‡], R Fu[§] and K Nasu[¶]

[†] Institute for Molecular Science, Myodaiji, Okazaki
444, Japan^{**}

[‡] National Lab for Infrared Physics of Academia Sinica
and Structure Research Lab, University of Science and
Technology of China, Hefei, China

X S Li, D L Lin and T F George

Department of Physics and Astronomy, State University of
New York at Buffalo, Buffalo, N.Y. 14260, USA

[¶] Permanent address: Department of Physics, Fudan Univer-
sity, Shanghai 200433, China

[§] Present address: Services de Chimie des Matériaux Non-
veaux, Université de l'Etat à Mons, B7000, Mons, Belgi-
que

^{**} Mailing address

Abstract

By considering the finite lifetime of excitation and the band structure of a finite polymer chain, the two-peak structure in the spectrum $\chi^{(3)}(\omega)$ of the third order nonlinear susceptibility of trans-polyacetylene is interpreted in this letter. The numerical result of our theoretical calculation agrees with the spectrum $\chi^{(3)}(\omega)$ observed in experiments.

In recent years significant progress has been achieved in the study of conducting polymers, where it is found that some conducting polymers possess not only unusually large third-order nonlinear optical susceptibility ($\chi^{(3)} > 10^{-9}$ esu) but also extremely fast response (sub-picosecond) [1-3]. Even in the infrared region far from the three-photon resonance, such organic material can still keep strong nonlinearity [4]. Noticing that the polymer has much smaller refractive index than the inorganic semiconductor, the effective nonlinear susceptibility $\chi^{(3)}/n^2$ of the polymer becomes more efficient. The advantages of both strong nonlinearity and quick response suggest that the conducting polymer is a promising material to produce ~~the~~ all-optical switching devices.

The spectrum $\chi^{(3)}(\omega)$ of the third-order nonlinear susceptibility of trans-polyacetylene has resonance struc-

true. By exploiting the resonance property, it is possible to enhance its nonlinearity further. The study of the spectrum structure is important for understanding and improving the optical nonlinear functions of the conducting polymer. The experiments demonstrate ^{that} there are two peaks in the spectrum $\chi''(\omega)$ of trans-polyacetylene, the main peak ~~fits~~ near 0.6 eV and the second one near 0.9 eV [1,3]. Since the optical gap 2Δ of trans-polyacetylene is about 1.8 eV, it is obvious that the position of the main peak is at one third of the gap and this peak is due to the three-photon resonance. Meanwhile the position of the second peak is at ~~the~~ half of the gap. As a naive speculation, it was thought that the second peak came from the two-photon resonance enhancement. But, in a careful consideration, it can be seen that the two-photon absorption in a rigid-band model is forbidden by momentum conservation [5]. So the second peak has some deep origin.

Recently many theoretical works have been devoted to explaining the structure of the third-order nonlinear susceptibility spectrum [5-10], but ~~this~~ ^{this} problem has not been solved yet. Actually this issue has become a hot topic in the area of conducting polymers. W. Wu and S. Kivelson established a systematic theory for the nonlinear susceptibility of the conducting polymer [10]. By using the Keldysh Green's Function method they derived an analytic expression for $\chi''(\omega)$ over the full frequency range. Their theoretical results rigorously present a divergence at $\hbar\omega = 2\Delta/3$ and a cusp at $\hbar\omega = \Delta$. The significance is χ''

there is ~~no~~ any adjustable parameter in their theory and the positions of the divergence and the cusp are in excellent agreement with those two peaks ^{observed in} of experiments. As is emphasized in their paper [10], the cusp is not the two-photon resonance enhancement. Due to the momentum conservation the transition matrix element for the two-photon process, vanishes at $\hbar\omega = \Delta$. They concluded that the second peak was induced by a large polarizability near that frequency.

However, this agreement between their theory and the experiments is challenged. C. Wu and one of the present authors (X.S.) ~~pointed out~~ ^{have pointed out} that their theory ~~was~~ ^{is} dealing with an ideal case, in which the damping of the excitation is neglected [11]. By using the Genkin-Mednis approach [12], C. Wu and X. Sun derive a general analytic formula of the nonlinear susceptibility from the SSH (Su-Schrieffer-Heeger) model [13]. In the limit of $\Delta/W \rightarrow 0$, where W is the bandwidth, C. Wu and Sun's formula of $\chi''(\omega)$ exactly reduces to that of W. Wu and S. Kivelson. It is to be expected, since W. Wu and Kivelson started from the TLM (Takahama-Lin-Liu-Maki) model [14], which is a continuum version of the SSH model. The point is, when a small damping of the excitation is taken into account, the cusp is greatly suppressed, and there is only a small bump left at that place [11]. The bump is too small to interpret the second peak near 0.9 eV. Due to the residual electron-lattice interaction and the imperfection of the lattice, the excitation always has some damping and its lifetime

should be finite. The cusp is so sensitive to the damping that a reasonable ^{by} small damping can almost eliminate it. So the origin of the second peak has to be studied further.

It should be noticed that in ~~the~~ realistic polymers not only the lifetime of the excitation but also the length of the chain are finite. The experiments have already shown that ~~the~~ polyenes possess strong chain-length dependence of the third-order nonlinear susceptibility [15]. The finiteness of the chain is also a substantial matter for the band structure of one-dimensional system. It is well known that the density of state in one-dimensional system has singularity at the band edges. But the divergence of the density of state appears only if the length of the chain is infinite. If the chain is not very long, e.g. in the case of trans-polyacetylene, the length is about 10^3 CH-monomers, the singularity will no longer persist, and the density of state inside the band is comparable with that at the band edges. Then the transition between the states inside the valence band and the conduction band, ~~which~~ ^{with} energy difference is larger than the gap 2Δ , will give considerable contribution to the third-order nonlinear susceptibility in the region with higher frequency than $\hbar\omega = 2\Delta/3$. As a result, the bump will be enhanced. The following calculation will show there indeed exists another distinct peak near 0.9eV when the length of the chain is shorter than 150 CH-monomers. It will also be seen ^{that} when the length of the chain is getting longer, the second peak will decrease. The reason is simple, when the length of the chain increases, the density

of states at the band edges will become larger and larger and the transition between the top edge of the valence band and the bottom edge of the conduction band dominant then only the main peak is left. It confirms the conclusion made in Ref. [11] that the infinite chain has only one peak at $\hbar\omega = 2\Delta/3$ after considering the damping of the excitation.

Now we are going to conduct the theoretical calculation in the light of the above analysis. Since the size of the chain is much smaller than the wavelength of the incident light, the chain can be considered as a big molecule, and the Orr and Ward's perturbation theory [16] can be used to calculate the third-order nonlinear susceptibility,

$$\chi_{\alpha\beta\gamma\delta}^{(3)}(-\omega; \omega_1, \omega_2, \omega_3) = \frac{K}{\hbar^3} \sum_{\alpha} \left\{ \sum_{m,n,p} \frac{\langle g | \mu_\alpha | m \rangle \langle m | \mu_\beta | n \rangle \langle n | \mu_\gamma | p \rangle \langle p | \mu_\delta | g \rangle}{(\omega_m - \omega)(\omega_n - \omega_1)(\omega_p - \omega_2)(\omega_g - \omega_3)} \right. \\ \left. - \sum_{m,n,p} \frac{\langle g | \mu_\alpha | m \rangle \langle m | \mu_\beta | p \rangle \langle p | \mu_\gamma | n \rangle \langle n | \mu_\delta | g \rangle}{(\omega_m - \omega)(\omega_n - \omega_1)(\omega_p + \omega_2)(\omega_g + \omega_3)} \right\}, \quad (1)$$

where $\omega = \omega_1 + \omega_2 + \omega_3$, $\alpha, \beta, \gamma, \delta$ are the component indices, g is ground state and m, n, p are excited states. K is a numerical factor, for the third harmonic generation, it is $1/4$. \sum_p is the summation for the permutation of the frequencies. μ_α is the α -component of the di-

pole. $\frac{1}{2}\omega_n$ is the energy difference between the excited state m and the ground state g .

For a finite dimerized chain consisting of $2N$ atoms and $2N$ electrons, the wave functions of both the ground state and the excited states can be obtained by solving the eigen-equation of the SSH Hamiltonian

$$H/t_0 = - \sum_{n,s} (1 + (-1)^n (\omega_{n+1} + \omega_n)) (a_{n,s}^\dagger a_{n+1,s} + h.c.) + \frac{1}{\pi\lambda} \sum_n (\omega_{n+1} + \omega_n)^2, \quad (2)$$

where t_0 is the hopping constant, λ the coupling constant, u_n the dimensionless displacement of lattice in site n , $a_{n,s}^\dagger$ and $a_{n,s}$ the creation and annihilation operators of electron in the site n with spin s . For the trans-polyacetylene

$$t_0 = 2.5\text{eV}, \quad \lambda = 0.2.$$

Since the photon momentum is very small, only the transitions with momentum conservation are important, and the dipole transition matrix between different excited states vanishes, then the Eq.(1) requires that the two-photon resonance is constrained, it is the same conclusion as W.Wu's [5].

Due to the strong anisotropy in the polymer chain, the direction along the chain is most important, and only the parallel component of the susceptibility is

needed to be calculated. For the third harmonic generation, $\omega_1 = \omega_2 = \omega$, then the spectrum $\chi^{(3)}(\omega)$ is

$$\chi^{(3)}(\omega) = \chi_{x,x,x,x}^{(3)}(-3\omega; \omega, \omega, \omega). \quad (3)$$

As it has been pointed that the excitation possesses damping, the excitation energy should include a small imaginary part $i\eta$, for the trans-polyacetylene, η is about 0.03 in the unit of 2Δ [8,9]. It is straightforward to make the numerical calculation for the spectrum $\chi^{(3)}(\omega)$ based on the above formulism.

Our theoretical results are shown in the Fig.1. The solid line is the spectrum of the chain with length $2N = 90$, it contains two distinct peaks near 0.6 eV and 0.9 eV respectively. This solid line is quite close to the experimental data, which are demonstrated in the Fig.1 by the crosses. In the Fig.1, a longer chain with $2N = 180$ is also presented by the dashed line. As we have analysed previously, when the length of the chain increases, the transition from the top edge of the valence band to the bottom edge of the conduction band gradually becomes dominant, then the second peak is getting weaker. So there appears a shoulder rather than a

peak near 0.9ev in the dashed line. It should be mentioned that the real material consists of chains with different lengths, it can be considered that the observed spectrum is some average over the different lengths. Evidently the results shown in the Fig.1 indicate that our theory is a reasonable interpretation for the spectrum $\chi^{(1)}(\omega)$ of the trans-polyacetylene. On the other hand, our theory has neglected many other effects, such as the electron correlation, the lattice fluctuation, various defects etc. These effects should be included further when we are going to improve our theory and to get better quantitative agreement with the experimental data.


XS wishes to thank the Institute for Molecular Science for the kind invitation as visiting professor and hospitality during his visit there. This investigation was partially supported by the Project 863, National Science Foundation of China and the US Office of Naval Research.

References

- [1] Kajzar F, Etemad S, Baker G L and Messler J 1987 Synth. Met. **17** 563
- [2] Heeger A J, Moses D and Sinclair M 1987 Synth. Met. **17** 343
- [3] Fann W S, Benson S, Wadey J M J, Etemad S, Baker G L and Kajzar F 1989 Phys. Rev. Lett. **62** 1492
- [4] Heeger A J, Kivelson S, Schrieffer J R and Su W P 1988 Rev. Mod. Phys. **60** 781
- [5] Wu W K 1988 Phys. Rev. Lett. **61** 1119
- [6] Heflin J R, Wong K Y, Zamani-Khamirli O and Garito A F 1988 Phys. Rev. B **38** 1573
- [7] deMeio C P and Silbey R 1988 J. Chem. Phys. **88** 2567
- [8] Mackie D M, Cohen R J and Glick A J 1989 Phys. Rev. B **39** 3442
- [9] Yu J, Friedman B, Baldwin P R and Su W P 1989 Phys. Rev. B **39** 12814
- [10] Wu W K and Kivelson S 1989 Synth. Met. **28** 575
- [11] Wu C and Sun X 1990 Phys. Rev. B **41** 12845
- [12] Genkin G M and Mednis P M 1968 Zh. Eksp. Teor. Fiz. **54** 1137
- [13] Su W P, Schrieffer J R and Heeger A J 1980 Phys. Rev. B **22** 2099
- [14] Takayama H, Lin-Liu Y R and Maki K 1980 Phys. Rev. B **21** 2388
- [15] Hermann J P and Duculing J 1974 J. Appl. Phys. **45** 5100
- [16] Orr B J and Ward J F 1971 Mol. Phys. **20** 513

Caption

Fig.1 The theoretical spectra $\chi''(\omega)$ of finite chains with $2N = 90$ (solid line) and $2N = 180$ (dashed line). The crosses are the experimental data.



Y90 Abstracts Distribution List for Solid State & Surface Chemistry

Professor John Baldeschwieler
Department of Chemistry
California Inst. of Technology
Pasadena, CA 91125

Professor John Eyler
Department of Chemistry
University of Florida
Gainesville, FL 32611

Dr. Sylvia Johnson
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025

Professor Paul Barbara
Department of Chemistry
University of Minnesota
Minneapolis, MN 55455-0431

Professor James Garvey
Department of Chemistry
State University of New York
Buffalo, NY 14214

Dr. Zakya Kafafi
Code 6551
Naval Research Laboratory
Washington, DC 20375-5000

Dr. Duncan Brown
Advanced Technology Materials
20-D Danbury Rd.
New Milford, CT 06776

Professor Steven George
Department of Chemistry
Stanford University
Stanford, CA 94305

Professor Larry Kesmodel
Department of Physics
Indiana University
Bloomington, IN 47403

Professor Stanley Bruckenstein
Department of Chemistry
State University of New York
Buffalo, NY 14214

Professor Tom George
Dept. of Chemistry and Physics
State University of New York
Buffalo, NY 14260

Professor Max Lagally
Dept. Metal. & Min. Engineering
University of Wisconsin
Madison, WI 53706

Professor Carolyn Cassady
Department of Chemistry
Miami University
Oxford, OH 45056

Dr. Robert Hamers
IBM T.J. Watson Research Center
P.O. Box 218
Yorktown Heights, N Y 10598

Dr. Stephen Lieberman
Code 522
Naval Ocean Systems Center
San Diego, CA 92152

Professor R.P.H. Chang
Dept. Matls. Sci. & Engineering
Northwestern University
Evanston, IL 60208

Professor Charles Harris
Department of Chemistry
University of California
Berkeley, CA 94720

Professor M.C. Lin
Department of Chemistry
Emory University
Atlanta, GA 30322

Professor Frank DiSalvo
Department of Chemistry
Cornell University
Ithaca, NY 14853

Professor John Hemminger
Department of Chemistry
University of California
Irvine, CA 92717

Professor Fred McLafferty
Department of Chemistry
Cornell University
Ithaca, NY 14853-1301

Dr. James Duncan
Federal Systems Division
Eastman Kodak Company
Rochester, NY 14650-2156

Professor Leonard Interrante
Department of Chemistry
Rensselaer Polytechnic Institute
Troy, NY 12181

Professor Horia Metiu
Department of Chemistry
University of California
Santa Barbara, CA 93106

Professor Arthur Ellis
Department of Chemistry
University of Wisconsin
Madison, WI 53706

Professor Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, NY 14853

Professor Larry Miller
Department of Chemistry
University of Minnesota
Minneapolis, MN 55455-0431

Professor Mustafa El-Sayed
Department of Chemistry
University of California
Los Angeles, CA 90024

Professor Eugene Irene
Department of Chemistry
University of North Carolina
Chapel Hill, NC 27514

Professor George Morrison
Department of Chemistry
Cornell University
Ithaca, NY 14853

Professor Daniel Neumark
Department of Chemistry
University of California
La Jolla, CA 92037

Professor Robert Whetten
Department of Chemistry
University of California
Los Angeles, CA 90024

Professor David Ramaker
Department of Chemistry
George Washington University
Washington, DC 20052

Professor R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, CA 90024

Gary Rubloff
I T.J. Watson Research Center
P.O. Box 218
Yorktown Heights, NY 10598

Professor Nicholas Winograd
Department of Chemistry
Pennsylvania State University
University Park, PA 16802

Professor Richard Smalley
Department of Chemistry
Rice University
P.O. Box 1892
Houston, TX 77251

Professor Aaron Wold
Department of Chemistry
Brown University
Providence, RI 02912

Professor Gerald Stringfellow
Dept. of Maths. Sci.
& Engineering
University of Utah
Salt Lake City, UT 84112

Professor Vicki Wysocki
Department of Chemistry
Virginia Commonwealth University
Richmond, VA 23284-2006

Professor Galen Stucky
Department of Chemistry
University of California
Santa Barbara, CA 93106

Professor John Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, PA 15260

Professor H. Tachikawa
Department of Chemistry
Michigan State University
East Lansing, MI 48824-1328

Professor William Unertl
Dept. for Surface Sci.
& Technology
University of Maine
Orono, ME 04469

Dr. Terrell Vanderah
Dept. of Chem. & Phys.
Naval Weapons Center
China Lake, CA 93555

Professor John Weaver
Dept. of Chem. Eng. & Mat. Sci.
University of Minnesota
Minneapolis, MN 55455

Professor Brad Weiner
Department of Chemistry
University of Puerto Rico
San Juan, Puerto Rico 00931